


Student Number: _____

Question 1 16 marks (Begin a new booklet) Marks



KAMBALA

Mathematics Extension 2

HSC Assessment Task 2

Half-Yearly Examination

March 2008

General Instructions

- Reading time – 5 minutes.
- Working time – 2 hours.
- Answer all questions in the writing booklets provided. Start each question in a new booklet.
- Board-approved calculators may be used.
- A table of standard integrals is provided.
- All necessary working should be shown in every question.

Total marks – 80

- Attempt Questions 1-5.
- All questions are of equal value.

- (a) Evaluate $|\sqrt{5} - 2i|$. 1
- (b) If $z = -3 - 4i$ find $\frac{1}{z}$ in the form $a + ib$. 2
- (c) Simplify $\frac{1+i^5}{1-i}$. 2
- (d) (i) Express $z = \frac{-1+i}{\sqrt{3+i}}$ in modulus-argument form. 2
- (ii) Hence evaluate $\cos \frac{7\pi}{12}$ in surd form. 2
- (e) Let ω be a non-real cube root of unity.
- (i) Show that $\omega^2 + \omega + 1 = 0$. 2
- (ii) Prove that $b^3 + c^3 = (b+c)(b+c\omega)(b+c\omega^2)$. 2
- (f) Find a polynomial $P(x)$ with real coefficients having $2i$ and $1 - 3i$ as zeroes. 3

Question 2 16 marks (Begin a new booklet)

Marks

- (a) Consider the function $f(x) = x - \ln(1+x^2)$. 3

Question 2 16 marks (Begin a new booklet)

Question 3 16 marks (Begin a new booklet)

Marks

(a) Find $\int \frac{x}{\sqrt{x+1}} dx$

2

(b) Solve for x : $\frac{x^2 - 5x}{4 - x} \leq -3$

3

(c) (i) Sketch the function $f(x) = x^2 - c^2$, where c is a positive constant, clearly indicating its vertex and intercepts.

1

(ii) Hence, without using calculus, draw separate sketches, at least $\frac{1}{3}$ of a page, for each of the following curves. Clearly indicate turning points.

(A) $f(x) = |x^2 - c^2|$

2

(B) $f(x) = \frac{1}{x^2 - c^2}$

2

(C) $f(x) = \sqrt{x^2 - c^2}$

2

(D) $f(x) = (x^2 - c^2)^2$

2

(E) $f(x) = (x^2 - c^2)^3$


2

(a) Consider the function $f(x) = x - \ln(1 + x^2)$.

Show that $f'(x) \geq 0$ for all values of x .

3

(b)



ABC is a triangle inscribed in a circle as shown above. The bisector of $\angle BAC$ meets BC in X and the circle at D .

(i) Prove that $\triangle ABX \sim \triangle ADC$

2

(ii) Prove that $AB \cdot AC = AD \cdot AX$

1

(iii) Prove that $AB \cdot AC = AX^2 + BX \cdot XC$

2

Question 3 continues next page

Question 3 continued

Question 4 (Begin a new booklet)

Marks

(c) A cubic equation has roots l, m and n . Given that

$$l + m + n = -3$$

$$l^2 + m^2 + n^2 = 29$$


$$lmn = -6$$

(i) By considering the expansion of $(l + m + n)^2$, show that the cubic equation is given by: 2

$$x^3 + 3x^2 - 10x + 6 = 0$$

(ii) Hence find the values of l, m and n . 3

(d) OACB is a rectangle where $OA = 2OB$. D is the point of intersection of the diagonals. The point B represents the complex number z .



Find in terms of z , the complex number represented by:

(i) A 1

(ii) D 2

(a) Assuming the result $\cos 3\theta = 4\cos^3\theta - 3\cos\theta$ and using a suitable substitution, solve the equation $8x^3 - 6x + 1 = 0$. 3

(b) (i) If x and y are real, prove that $x^2 + y^2 \geq 2xy$. 2

(ii) Hence show that $a^2 + b^2 + c^2 + d^2 \geq 4abcd$. 2

(c) $P(2ap, ap^2)$ is a point on the parabola $x^2 = 4ay$. F is the focus of the parabola. PQ is the perpendicular from P to the directrix d , of the parabola. The tangent at P to the parabola, cuts the axis of the parabola at the point R . 2

(i) Show that the tangent at the point P to the parabola has equation

$$px - y - ap^2 = 0$$

(ii) Show that PR and QF bisect each other. 3

(iii) Show that $PR \perp QF$. 2

(iv) What type of quadrilateral is $PQRF$? Give reasons for your answer. 2

Question 5 continued

Marks

Question 5 (Begin a new booklet)

Marks

(b) For all integers $n \geq 1, I_n = \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n-1} + \frac{1}{2n}$.

Question 5 16 marks (Begin a new booklet) Marks

(a) Let $y = uv$ be the product of u and v , where u and v are functions of x .

(i) Show that $y'' = uv'' + 2u'v' + u''v$. 2

(ii) Find similar expressions for $y''', y^{(4)}$ and $y^{(5)}$. 2

(iii) Hence or otherwise, find and simplify $\frac{d^5}{dx^5} \left((1-x^2)e^{-x} \right)$. 2


Question 5 continues next page

Question 5 continued Marks

(b) For all integers $n \geq 1$, $t_n = \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n-1} + \frac{1}{2n}$.

(i) Show that $t_n + \frac{1}{2n} = \frac{1}{n} + \frac{1}{n+1} + \dots + \frac{1}{2n-1}$ 1

(ii)



The diagram above shows the graph of the function $y = \frac{1}{x}$ for $n \leq x \leq 2n$.

Using the diagram, show that $t_n + \frac{1}{2n} > \ln 2$. 3

(iii) For all integers $n \geq 1$, let $s_n = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{1}{2n-1} - \frac{1}{2n}$.

Using Mathematical Induction, prove that $s_n = t_n$. 4

(iv) Hence find, to three decimal places, the value of:

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{1}{9999} - \frac{1}{10000}$$
 2

End of Examination

Qn	Solutions	Marks	Comments
	<u>Kambala Extn 2 Half-Yearly Exam 2008</u>		
	<u>Question 1</u>		
(a)	$ 5-2i = \sqrt{5^2+2^2} = \sqrt{29}$	1	
(b)	$z = -3-4i$ $\frac{1}{z} = \frac{1}{-3-4i} \times \frac{-3+4i}{-3+4i}$ $= \frac{-3+4i}{-9+16}$ $= -\frac{3}{25} + \frac{4}{25}i$	1	
(c)	$\frac{1+i}{1-i}$ $= \frac{1+i}{1-i} \times \frac{1+i}{1+i}$ $= \frac{1+2i+1}{1+1}$ $= \frac{2+2i}{2} = 1+i$ $i^4 = 1$ $i^5 = i$	1	
(d)	(i) $z = \frac{-1+i}{\sqrt{3}+i}$ $(-1+i) = \sqrt{2} \operatorname{cis} \frac{3\pi}{4}$ $(\sqrt{3}+i) = 2 \operatorname{cis} \frac{\pi}{6}$ $z = \frac{\sqrt{2} \operatorname{cis} \frac{3\pi}{4}}{2 \operatorname{cis} \frac{\pi}{6}}$ $= \frac{\sqrt{2}}{2} \operatorname{cis} \left(\frac{3\pi}{4} - \frac{\pi}{6} \right)$ $= \frac{\sqrt{2}}{2} \operatorname{cis} \left(\frac{7\pi}{12} \right)$ (ii) $\frac{\sqrt{2}}{2} \left(\cos \frac{7\pi}{12} + i \sin \frac{7\pi}{12} \right) = \frac{-1+i}{\sqrt{3}+i}$ $\frac{-1+i}{\sqrt{3}+i} \times \frac{\sqrt{3}-i}{\sqrt{3}-i} = \frac{-\sqrt{3}+i+\sqrt{3}i+1}{3+1} = \frac{-\sqrt{3}i+i+\sqrt{3}+1}{4}$ Equating real parts $\frac{\sqrt{2}}{2} \cos \frac{7\pi}{12} = \frac{1+\sqrt{3}}{4}$ $\cos \frac{7\pi}{12} = \frac{1+\sqrt{3}}{2\sqrt{2}}$	1	

Qn	Solutions	Marks	Comments
(f)	<u>Question 1 ctd.</u> $P(z) = (z-2i)(z-(1-3i))$ $(z-2i)(z-1+3i)$		

Qn	Solutions	Marks	Comments
	<u>Question 1 ctd.</u>		
(e)	(i) w is a non-real cube root of unity, <u>Method I</u> $w^3 = 1$ $w^3 - 1 = 0$ $(w-1)(w^2+w+1) = 0$ $\therefore w = 1$ or $w^2+w+1 = 0$ but w is a non-real cube root of unity $w \neq 1$ so $w^2+w+1 = 0$	1	
	<u>Method II</u> $z^3 = 1$ Let $z = 1 \operatorname{cis} \theta$ $\operatorname{cis} 3\theta = \operatorname{cis} 0$ $3\theta = 0 + 2k\pi$ $\theta = \frac{2k\pi}{3}$ When $k=0$ $z_1 = \operatorname{cis} 0 = 1$ $k=1$ $z_2 = \operatorname{cis} \frac{2\pi}{3} = w$ $k=2$ $z_3 = \operatorname{cis} \frac{4\pi}{3} = w^2$ For $z^3 - 1 = 0$ Sum roots = 0 $\therefore 1 + w + w^2 = 0$	1	
	<u>Method III</u> $w^3 = 1$ $(w^2)^3 = (w^3)^2 = 1$ $\therefore w^2$ is also a root $1^3 = 1$ and 1 is obviously a root $\therefore w^2, w$ and 1 are cube roots of unity For $z^3 - 1 = 0$ Sum of roots = 0 $\therefore w^2 + w + 1 = 0$	1	
(ii)	$w^2 + w + 1 = 0$ RHS = $(b+ic)(b+cw)(b+cw^2)$ $w^3 = 1$ $= (b+ic)(b^2+bcw^2+bcw+c^2w^3)$ $w^4+w+1=0$ $= (b+ic)(b^2-bc+ic^2)$ $\therefore w^2+w = -1$ $= b^3+ic^3 = \text{LHS} \quad \sin a x^2 + 4^3 = (x+4)(x^2-2x+4)$ <u>Method III (Similar)</u> $(b^2-bc+ic^2)$ LHS $b^3+ic^3 = (b+ic)(b+cw)(b+cw^2)$ RHS = $(b+ic)(b+cw)(b+cw^2)$ $(b+cw)(b+cw^2) = b^2+bcw^2+bcw+c^2w^3$ $= b^2+bc(w^2+w)+c^2$ $= b^2-bc+ic^2$	1	

Qn	Solutions	Marks	Comments+Criteria
2 (a)	$\int \frac{x \, dx}{x^2+1}$		

$$(b^2+c^2)(b^2+c^2) = b^2 + bcw^2 + bcw^2 + c^2w^2$$

$$= b^2 + bc(w^2+w) + c^2$$

$$= b^2 - bc + c^2$$




$\therefore LHS = RHS$

$$\cos \frac{\pi}{12} = \frac{\sqrt{3}}{4} \times \frac{\sqrt{2}}{\sqrt{2}} + \frac{\sqrt{2}}{4} \times \frac{\sqrt{3}}{\sqrt{2}}$$




$$= \frac{\sqrt{2}(\sqrt{3} + \sqrt{3})}{4} \text{ or } \frac{\sqrt{2} \cdot 2\sqrt{3}}{4}$$

Qn	Solutions	Marks	Comments
(f)	<p>Question 1 ctd.</p> $P(z) = (z-2i)(z-(1-3i))$ $(z+2i)(z-(1+3i))$ <p>If $z = 2i$ and $1-3i$ are zeros then $-2i$ and $1+3i$ are also zeros as $P(z)$ has real coefficients.</p> $(z-2i)(z+2i) = z^2 + 4$ $(z-(1-3i))(z-(1+3i))$ $= z^2 - (1-3i+1+3i)z + 1+9$ $= z^2 - 2z + 10$ $P(z) = (z^2+4)(z^2-2z+10)$ $= z^4 - 2z^3 + 10z^2 + 4z^2 - 8z + 40$ $= z^4 - 2z^3 + 14z^2 - 8z + 40$ <p>$\therefore P(z)$ could be</p> $P(z) = z^4 - 2z^3 + 14z^2 - 8z + 40$	1	

Qn	Solutions	Marks	Comments+Criteria
2 (a)	$\int \frac{x \, dx}{\sqrt{x+1}}$ <p>Let $u = x+1$ $\therefore du = dx$ and $x = u-1$</p> $\int \frac{x}{\sqrt{x+1}} \, dx$ $= \int \frac{u-1}{\sqrt{u}} \cdot du$ $= \int \frac{u}{\sqrt{u}} - \frac{1}{\sqrt{u}} \, du$ $= \int u^{\frac{1}{2}} - u^{-\frac{1}{2}} \, du$ $= \frac{2u^{\frac{3}{2}}}{\frac{3}{2}} - 2u^{\frac{1}{2}} + C$ $= \frac{2}{3} \sqrt{x+1}^3 + 2\sqrt{x+1} + C$		
(b)	$x^2 - 5x \leq -3$ $\frac{x^2 - 5x}{4-x} \leq -3(4-x)^2$ $x(x-5)(4-x) + 3(4-x)^2 \leq 0$ $(4-x)(x(x-5) + 3(4-x)) \leq 0$ $(4-x)(x^2 - 5x + 12 - 3x) \leq 0$ $(4-x)(x^2 - 8x + 12) \leq 0$ $(4-x)(x-6)(x-2) \leq 0$ <p>$x = 2, 6, 4$ if $x=0, y=4-6 \cdot 2$</p> <p>$\therefore -2 \leq x < 4$ and $x > 6$</p>		


Qn	Solutions	Marks	Comments+Criteria
2 c) d)	<p>(i) $f(x) = x^2 - c^2, c > 0$</p> 		
	<p>(ii) (A) $f(x) = x^2 - c^2$</p> 		
	<p>(B) $f(x) = \frac{1}{x^2 - c^2} = \frac{1}{(x-c)(x+c)}$</p> <p>$x \neq -c$</p>  <p>if $x=0, f(x) = \frac{1}{-c^2}$</p>		

Qn	Solutions	Marks	Comments+Criteria
2 c) d)			

Qn	Solutions	Marks	Comments+Criteria
2 c) d)	<p>(c) c) d)</p> <p>(c) $f(x) = \sqrt{x^2 - c^2}$</p> 		
	<p>(D) $f(x) = (x^2 - c^2)^2$</p> <p>$= (x^2 - c^2)(x^2 - c^2)$</p> <p>$= (x - c)^2 (x + c)^2$</p> 		
	<p>(E) $f(x) = (x^2 - c^2)^3$</p> <p>$= (x - c)^3 (x + c)^3$</p> 		

Qn	Solutions	Marks	Comments+Criteria
3 c) d)			

Qn	Solutions	Marks	Comments+Criteria
3 Sol	<p>(i) Sol</p> <p>(ii) Prove $AB \cdot AC = AD \cdot AX$</p> <p>Since $\triangle ABX \parallel \triangle ADC$ from (i), then corresponding sides are in proportion.</p> $\therefore \frac{AB}{AX} = \frac{AD}{AC}$ $\therefore AB \cdot AC = AD \cdot AX$ <p>(iii) Prove $AB \cdot AC = AX^2 + BX \cdot XC$</p> <p>From (i), $AB \cdot AC = AD \cdot AX$</p> $= (AX + XD) \cdot AX$ $= AX^2 + XD \cdot AX$ $= AX^2 + BX \cdot XC$ <p>since $AX \cdot XD = BX \cdot XC$ by intercept theorem</p>		

Qn	Solutions	Marks	Comments+Criteria
3	<p>(a) $f(x) = x - \ln(1+x^2)$</p> $f'(x) = 1 - \frac{2x}{1+x^2}$ $= \frac{1+x^2-2x}{1+x^2}$ $= \frac{x^2-2x+1}{1+x^2}$ $= \frac{(x-1)^2}{1+x^2}$ <p>Since $x^2 > 0$, then $(x-1)^2 > 0$</p> <p>$(x-1)^2 > 0 \therefore f'(x) > 0 \forall x$</p>		
(b)	 <p>(1) Prove $\triangle ABX \parallel \triangle ADC$</p> <p>$\therefore \triangle ABX$ and $\triangle ADC$</p> <p>AD is bisector of $\angle BAC$ (data)</p> <p>$\therefore \angle CAX = \angle BAX$</p> <p>$\angle AXC = \angle ABX$ (angles in same arc =)</p> <p>$\therefore \angle ACX = \angle AXB$ (angle sum of \triangle)</p> <p>$\therefore \triangle ABX \parallel \triangle ADC$ (equiangular)</p>		


Qn	Solutions	Marks	Comments+Criteria
3	<p>(c) $l+mn = -3 = -2x$ cubic $l^2 + m^2 + n^2 = 29$ $l^2 + m^2 = ?$ $lmn = -6$ $l^2 + m^2 = ?$</p> <p>(i) $(l+mn)^2$ $= l^2 + m^2 + n^2 + 2lm + 2ln + 2mn$ $= l^2 + m^2 + n^2 + 2(lm + mn + nl)$</p> <p>$(-3)^2 = 29 + 2(lm + mn + nl)$ $9 = 29 + 2(lm + mn + nl)$ $lm + mn + nl = -\frac{20}{2}$ $= -10$</p> <p>cubic is given by $x^3 - (sum)x^2 + (product of 2)x - product = 0$</p> <p>$\therefore x^3 + 3x^2 - 10x + 6 = 0$ as reqd</p> <p>(ii) $x^3 + 3x^2 - 10x + 6 = 0 = P(x)$ $P(1) = 0$. $\therefore x=1$ is a factor.</p> $\begin{array}{r} x^3 + 3x^2 - 10x + 6 \\ \underline{x^3 - x^2} \\ 4x^2 - 10x + 6 \\ \underline{4x^2 - 4x} \\ -6x + 6 \\ \underline{-6x + 6} \\ 0 \end{array}$		

Qn	Solutions	Marks	Comments
(a)	<p>Question 4</p> <p>$\cos 3\theta = 4\cos^3\theta - 3\cos\theta$</p>		

Qn	Solutions	Marks	Comments+Criteria
3	<p>(c) det $\therefore P(x) = (x-1)(x^2 + 4x - 6)$ $x=1$ a root, $x = -4 \pm \sqrt{16 - 4 \cdot 1 \cdot (-6)}$</p> $\begin{array}{l} \frac{2 \cdot 1}{2} \\ = -4 \pm \frac{\sqrt{16 + 24}}{2} \\ = -4 \pm \frac{\sqrt{40}}{2} \\ = -4 \pm \frac{2\sqrt{10}}{2} \\ x = -2 \pm \sqrt{10} \end{array}$ <p>$\therefore 4, m, n$ are $1, -2 \pm \sqrt{10}$.</p>		

Qn	Solutions	Marks	Comments
(a)	<p>Question 4</p> <p>$\cos 3\theta = 4\cos^3\theta - 3\cos\theta$</p>		

$16x + 6$



(i) let $z = be^{-i\theta} (\cos \theta + i \sin \theta)$

$OB = z$

$OA = zOB$

$OA = z^2 = 2z$ (rotation anticlockwise by $\frac{\pi}{2}$)

$\therefore A$ is $2iz$

(ii) $OC = OB + BC$

$= z + 2iz$

$= (1+2i)z$

$\therefore OB = \frac{1}{2}(1+2i)z$

$\therefore D$ is $\frac{1}{2}(1+2i)z$

or $(\frac{1}{2} + i)z$

OR SOLUTIONS

Question 4

$\cos 3\theta = 4\cos^3 \theta - 3\cos \theta$

(a) $8x^3 - 6x + 1 = 0$

$8x^3 - 6x = -1$

$4x^3 - 3x = -\frac{1}{2}$

looking for 3 solutions (unique), let $x = \cos \theta$

$4\cos^3 \theta - 3\cos \theta = -\frac{1}{2}$

$\cos 3\theta = -\frac{1}{2}$

$3\theta = \pi - \frac{\pi}{3}, \pi - \frac{2\pi}{3}$

$= \frac{2\pi}{3} + 2k\pi, -\frac{2\pi}{3} + 2k\pi$

$3\theta = \frac{6k\pi \pm 2\pi}{3} = 2\pi(3k \pm 1)$

$\theta = \frac{2\pi}{3}(3k \pm 1)$

when $k=0$ $\theta = \frac{2\pi}{3} \therefore x = \cos \frac{2\pi}{3}$

when $k=1$ $\theta = \frac{2\pi \times 2}{3} = \frac{4\pi}{3} \therefore x = \cos \frac{4\pi}{3}$

$\theta = \frac{2\pi \times 2}{3} = \frac{4\pi}{3} \therefore x = \cos \frac{4\pi}{3}$

when $k=2$ $\theta = \frac{2\pi \times 7}{3} = \frac{14\pi}{3} = \frac{2\pi}{3} \therefore x = \cos \frac{2\pi}{3}$

$\theta = \frac{2\pi \times 5}{3} = \frac{10\pi}{3} = \frac{4\pi}{3} \therefore x = \cos \frac{4\pi}{3}$

$\theta = \frac{2\pi \times 5}{3} = \frac{10\pi}{3} = \frac{4\pi}{3} \therefore x = \cos \frac{4\pi}{3}$

$\therefore x = \cos \frac{2\pi}{3}, \cos \frac{4\pi}{3}, \cos \frac{14\pi}{3}$

$\cos \frac{14\pi}{3} = \cos \frac{2\pi}{3}$

Qn	Solutions	Marks	Comments+Criteria
4	<p>(b) (i) have $x^2 + y^2 > 2xy$ Now $(x-y)^2 > 0 \quad \forall x, y$ real $\therefore x^2 - 2xy + y^2 > 0$ $\Rightarrow x^2 + y^2 > 2xy$</p>		

Q4(b) see over

4			
---	--	--	--

Qn	Solutions	Marks	Comments+Criteria
4	<p>(c) dtd. (i) $P(2ap, ap^2)$ $y = \frac{1}{4a}x^2$ $\frac{dy}{dx} = \frac{1}{2a}x$ $m = \frac{2ap}{2a} = p$ $y - ap^2 = p(x - 2ap)$ $y - ap^2 = px - 2ap^2$ $\therefore px - ap^2 - y = 0$</p> <p>(ii) Show PR and OF bisect. R lies on tangent and axis of parabola. \therefore at $x=0, 0 - ap^2 - y = 0$ $\therefore y = -ap^2$ R is $(0, -ap^2)$. midpoint of PR = $(\frac{2ap+0}{2}, \frac{-ap^2+ap^2}{2})$ $= (ap, 0)$ midpoint of OF = $(\frac{2ap+0}{2}, \frac{-a+a}{2})$ $= (ap, 0)$ since midpoint of PR = midpoint of OF Then PR and OF must bisect</p>		

Qn	Solutions	Marks	Comments+Criteria
4	(c) dtd		

Qn	Solutions	Marks	Comments+Criteria
5	(a) $y = uv$		

Qn	Solutions	Marks	Comments+Criteria
4	<p>(e) dtd</p> <p>(ii) Show PR \perp OF</p> $M_{PR} = \frac{ap^2 + ap^2}{2ap - 0}$ $= \frac{2ap^2}{2ap}$ $= p \quad (\text{already shown in (i)!})$ $M_{OF} = \frac{-a - a}{2ap - 0}$ $= \frac{-2a}{2ap}$ $= -\frac{1}{p}$ <p>$M_{PR} \times M_{OF} = p \times -\frac{1}{p}$</p> $= -1$ <p>$\therefore PR \perp OF$</p> <p>(iv) PORF?</p> <p>Diagonals meet @ right angles. \therefore PORF is a rhombus.</p> <p>FR \parallel PO since PR is perp. to diagonal, a horizontal line.</p>		

Qn	Solutions	Marks	Comments+Criteria
5	<p>(a) $y = uv$</p> <p>(i) $y' = u'v + v'u$</p> $y'' = u''v + v'u' + v''u + u'v'$ $= uv'' + 2u'v' + u''v$ <p>(ii) find y''', y''''</p> $y''' = u'v'' + v'''u + 2u''v' + v''u'' + u''v' + v'u''$ $= uv''' + 3u'v'' + 3u''v' + u'''v$ $y'''' = u'v'''' + v''''u + 3u''v''' + v''''u' + 3u''v'' + u''''v + v''''u'' + 6u'v'''' + 4u''v''' + 4u''v'' + u''''v$ $y'''' = u'v'''' + v''''u + 6u''v''' + v''''u' + 4u''v'' + v''''u'' + 4u''v'' + 5u'v'''' + 10u''v''' + 10u''v'' + 6u''v'' + v''''u'' + 4u''v'' + v''''u'' + 4u''v'' + 5u''v'' + u''''v$		
(ii)	$\frac{d^2}{dx^2} \left(\frac{(1-x^2)e^{-x}}{u} \right)$ $u = 1-x^2$ $\frac{du}{dx} = u' = -2x$ $y'' = uv'' + 5u'v' + 10u''v + 10u''v' + 5u''v'' + u''v''$ $\therefore \frac{d^5}{dx^5} \left((1-x^2)e^{-x} \right)$ $v = e^{-x}$ $v' = -e^{-x}$ $v'' = e^{-x}$ $v''' = -e^{-x}$ $v'''' = e^{-x}$ $v'''' = -e^{-x}$ $= (1-x^2)(-e^{-x}) + 5(-2x)e^{-x} + 10(-2)(-e^{-x})$ $+ 10 \cdot 0 \dots + 5 \times 10 \dots + 0$ $= -e^{-x}(1-x^2 + 10x - 20)$ $= -e^{-x}(-x^2 + 10x - 19) = e^{-x}(x^2 - 10x + 19)$		

Qn	Solutions	Marks	Comments+Criteria
5	$t_n = \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n-1} + \frac{1}{2n}$ $(i) t_n + \frac{1}{2n} = \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n-1} + \frac{1}{2n} + \frac{1}{2n}$ $= \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n-1} + \frac{2}{2n}$ $= \frac{1}{n} + \frac{1}{n+1} + \dots + \frac{1}{2n-1}$ <p style="text-align: center;">as required</p> $(ii) \int_n^{2n} \frac{1}{x} dx \approx n \times \frac{1}{n} + n \times \frac{1}{n+1} + \dots + n \times \frac{1}{2n-1} + n \times \frac{1}{2n}$ $= 1 \cdot \left(\frac{1}{n} + \frac{1}{n+1} + \dots + \frac{1}{2n-1} \right) = \left(t_n + \frac{1}{2n} \right)$ <p>also $\int_n^{2n} \frac{1}{x} dx = \ln 2n - \ln n = \ln 2n - \ln n = \ln 2 + \ln n - \ln n = \ln 2$</p> <p>From diagram, area of rectangle is an approximation: giving an area greater than exact area.</p> $\therefore \left(\frac{1}{n} + \frac{1}{n+1} + \dots + \frac{1}{2n-1} \right) > \ln 2$ $\text{i.e. } \left(t_n + \frac{1}{2n} \right) > \ln 2$		

Qn	Solutions	Marks	Comments+Criteria
5	$(iii) S_n = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{1}{2n-1} - \frac{1}{2n}$ <p>Prove $S_n = t_n$.</p> $t_n = \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n-1} + \frac{1}{2n}$ <p>RTP:</p> $1 - \frac{1}{2} + \frac{1}{3} - \dots + \frac{1}{2n-1} - \frac{1}{2n} = \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n-1} + \frac{1}{2n} \text{ for } n > 1.$ <p>Let $n=1$:</p> $\text{LHS} = 1 - \frac{1}{2} = \frac{1}{2}$ $\text{RHS} = \frac{1}{1+1} = \frac{1}{2} = \text{LHS}$ <p>\therefore true for $n=1$.</p> <p>Assume true for $n=k$ i.e. $S_k = t_k$.</p> $\text{i.e. } 1 - \frac{1}{2} + \frac{1}{3} - \dots + \frac{1}{2k-1} - \frac{1}{2k} = \frac{1}{k+1} + \frac{1}{k+2} + \dots + \frac{1}{2k-1} + \frac{1}{2k}$ <p>Prove true for $n=k+1$ i.e. $S_{k+1} = t_{k+1}$.</p> <p>LTP:</p> $1 - \frac{1}{2} + \frac{1}{3} - \dots + \frac{1}{2(k+1)-1} - \frac{1}{2(k+1)} = \frac{1}{(k+1)+1} + \frac{1}{(k+1)+2} + \dots + \frac{1}{2k+1} - \frac{1}{2k+2}$ $\text{i.e. } 1 - \frac{1}{2} + \frac{1}{3} - \dots + \frac{1}{2k+1} - \frac{1}{2k+2} = \frac{1}{k+2} + \frac{1}{k+3} + \dots + \frac{1}{2k+1} + \frac{1}{2k+2}$ $\text{LHS} = 1 - \frac{1}{2} + \frac{1}{3} - \dots + \frac{1}{2k+1} - \frac{1}{2k+2}$ $= 1 - \frac{1}{2} + \frac{1}{3} - \dots - \frac{1}{2k} + \frac{1}{2k+1} - \frac{1}{2k+2}$ $= \frac{1}{k+1} + \frac{1}{k+2} + \dots + \frac{1}{2k-1} + \frac{1}{2k} + \frac{1}{2k+1} - \frac{1}{2k+2}$ $= \frac{1}{k+2} + \frac{1}{k+3} + \dots + \frac{1}{2k+1} + \frac{1}{2k+2} - \frac{1}{2k+2}$ $= \frac{1}{k+2} + \frac{1}{k+3} + \dots + \frac{1}{2k+1} + \frac{1}{2k+2} - \frac{1}{2k+2}$ $= \frac{1}{k+2} + \frac{1}{k+3} + \dots + \frac{1}{2k+1} + \frac{1}{2k+2}$		

Qn	Solutions	Marks	Comments+Criteria
5	$(iv) 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{1}{999} - \frac{1}{1000}$		

Qn	Solution	Marks	Comments+Criteria
5 (iv)	$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{1}{9999} - \frac{1}{10000}$		
	$S_n = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{1}{2n-1} - \frac{1}{2n}$		
	<p>So $2n = 10000$ $n = 5000$</p>		
	$t_n = S_n \therefore t_{5000} = S_{5000}$		
	$t_n + \frac{1}{2n} > \ln 2$		
	$t_{5000} + \frac{1}{10000} > \ln 2$		
	$t_{5000} > \ln 2 - \frac{1}{10000}$		
	$\therefore t_{5000} \doteq 0.693$		

$$= \frac{1}{k+2} + \frac{1}{k+3} + \dots + \frac{1}{2k+1} + \frac{1}{2k+2}$$

$$= \frac{1}{2k+2} + \frac{1}{2k+3} + \dots + \frac{1}{2k+1} + \frac{1}{2k+2}$$